

Andrex

APPEARANCE:

Leaflet

COLOR KEY

■ Process BLACK

FONTS:

Arial Regular/Bold/Italic

DIMENSIONS:


In Millimeters (mm). Deviation + 0.5 mm

Width

Height

95.25

152.4



OBS
Andrex Capsules

(Testosterone Undecanoate)

For oral use only

Composition

Each capsule contains:
40 mg testosterone undecanoate (B.P.) dissolved in oleic acid.

Characteristics

Andrex is an orally active testosterone preparation. The active substance is testosterone undecanoate, a fatty acid ester of the natural androgen testosterone.

Testosterone as such is inactive on oral administration. Testosterone undecanoate is able to by pass the liver via the lymphatic system and is therefore orally active.

Therapy with **Andrex** increases plasma levels of testosterone and its active metabolites, leading to a regular therapeutic effect.

Andrex is well tolerated and unlike the orally active C-17-methylated testosterone derivatives, does not affect liver function.

Indications

Testosterone replacement therapy in male hypogonadal disorders for example:

- after castration
- eunuchoidism
- hypopituitarism
- endocrine impotence
- male climacteric symptoms such as decreased libido and decreased mental and physical activity
- certain types of infertility due to disorders of spermatogenesis

Moreover, testosterone therapy may be indicated in osteoporosis due to androgen deficiency.

خوراک: ڈاکٹر کے مشورے کے مطابق استعمال کریں۔

Dosage

In general, dosage should be adjusted according to the response of the individual patient. Usually, an initial dosage of 120-160 mg daily for 2-3 weeks is adequate, followed by a maintenance dosage of 40-120 mg daily.

اینڈریکس کیپسولز
(ٹیسٹوسٹیرون ان ڈیکانویٹ)
صرف کھانے کے لئے

Administration

Andrex capsules should be taken after meals, with a little fluid if necessary and swallowed whole without chewing. It is preferable that half of the daily dose be taken in the morning and the other half in the evening. If an uneven number of capsules is to be taken, the larger dose should be taken in the morning.

Contra-indications

Known or suspected prostatic or mammary carcinoma

Warnings and precautions

- If androgen associated adverse reactions occur, treatment should be interrupted and, after disappearance of the symptoms, be resumed at a lower dosage.
- Patients with latent or overt cardiac failure, renal dysfunction, hypertension, epilepsy or migraine (or a history of these conditions) should be monitored since, androgens may occasionally induce salt and fluid retention.
- Androgens should be used cautiously in prepubertal boys to avoid premature epiphyseal closure or precocious sexual development.
- A decrease in protein-bound iodine (PBI) may occur, but this has no clinical significance.

Adverse reactions

The following adverse reactions have been associated with androgen therapy:

- Priapism and other signs of excessive sexual stimulation.
- In prepubertal boys, precocious sexual development, an increased frequency of erections, phallic enlargement and premature epiphyseal closure.
- Oligospermia and decreased ejaculatory volume.
- Water & salt retention.

Presentation

3x10's Strips

Instructions

Store at 2-8°C. Protect from sunlight & moisture.

Avoid freezing. Keep out of reach of children.

M.L. No. : 000012
Manufactured By:
OBS Pakistan (Pvt) Ltd.
C-14, S.I.T.E., Karachi-75700

CIR-AND-02/OBS/10

ہدایات: ۸۲۲ ڈگری سینٹی گریڈ درجہ حرارت کے درمیان رکھیں۔
دھوپ اور نمی سے بچائیں۔ ٹھنڈ ہونے سے بچائیں۔ بچوں کی پہنچ سے دور رکھیں۔

OBS

Mfg. Specs. OBS.

PACKAGING COMPONENTS APPROVAL

MARKETING	
REGULATORY	
TECH. OPERATION	
SUPPLY CHAIN	